

Early-Time Chromatic Variations in the Wind-Swept Medium of GRB 021211 and the Faintness of its Afterglow

M. C. Nysewander¹, D. E. Reichart¹, H.-S. Park², G. G. Williams³, K. Kinugasa⁴, D. Q. Lamb⁵, A. A. Henden⁶, S. Klose⁷, T. Kato⁸, A. Harper⁵, H. Yamaoka⁹, C. Laws¹⁰, K. Torii¹¹, D. G. York⁵, J. C. Barentine¹², J. Dembicky¹², R. J. McMillan¹², J. A. Moran¹, D. H. Hartmann¹³, B. Ketzeback¹², M. B. Bayliss¹, J. W. Bartelme¹, J. A. Crain¹, A. C. Foster¹, M. Schwartz¹⁴, P. Holvorcem¹⁴, P. A. Price¹⁵, R. Canterna¹⁶, G. B. Crew¹⁷, G. R. Ricker¹⁷, and S. D. Barthelmy¹⁸

¹Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Campus Box 3255, Chapel Hill, NC 27599; mnysewan@physics.unc.edu, reichart@physics.unc.edu

²Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550

³MMT Observatory, University of Arizona, Tucson, AZ 85721

⁴Gunma Astronomical Observatory, 6860-86 Nakayama Takayama, Agatsuma, Gunma 377-0702, Japan

⁵Department of Astronomy & Astrophysics, University of Chicago, 5640 S. Ellis Ave., Chicago, IL, 60615

⁶AAVSO, Clinton B. Ford Astronomical Data and Research Center, 25 Birch St., Cambridge, MA

⁷Thüringer Landessternwarte, 07778 Tautenburg, Germany

⁸Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan

⁹Department of Physics, Kyushu University, Fukuoka 810-8560, Japan

¹⁰Department of Astronomy, Box 351580, University of Washington, Seattle, WA 98195

¹¹Department of Earth and Space Science, Graduate School of Science, 1-1 Machikaneyama-cho, Toyonaka, Osaka 560-0043, Japan

¹²Apache Point Observatory, P.O. Box 59, Sunspot, NM 88349

¹³Clemson University, Department of Physics and Astronomy, Clemson, SC 29634

¹⁴Tenagra Observatory, HC2 Box 292, Nogales, AZ 85621

¹⁵University of Hawaii, Institute of Astronomy, 2680 Woodlawn Drive, Honolulu, HI 96822-1897

¹⁶University of Wyoming, Department of Physics and Astronomy, P.O. Box 3905, Laramie, WY 82072

¹⁷Center for Space Research, Massachusetts Institute of Technology, 70 Vassar Street, Cambridge, MA, 02139

¹⁸NASA Goddard Space Flight Center, Code 661, Greenbelt, MD 20771

ABSTRACT

We present Follow-Up Network for Gamma-Ray Bursts (FUN GRB) Collaboration observations of the optical afterglow of GRB 021211 made between 143 seconds and 102 days after the burst. Our unique data set includes the earliest filtered detections and color information for an afterglow in the pre-Swift era. We find that the afterglow is best described by (1) propagation through a wind-swept medium, (2) a cooling break that is blueward of the observed optical frequencies, and (3) a hard electron energy distribution. However, superimposed on this “standard model” behavior we find one and possibly two significant chromatic variations during the first few hours after the burst. We consider possible reasons for these variations, including the possibility that they are due to a dust echo. Finally, we constrain physical parameters that describe the afterglow and surrounding medium for a variety of scenarios and find that GRB 021211’s afterglow is faint for a combination of 3 – 4 reasons: (1) a low fraction of energy in relativistic electrons, (2) a low density for the wind-swept medium, implying either a low mass-loss rate and/or a high wind velocity for the progenitor, (3) a wide opening/viewing angle for the jet, and possibly (4) moderate source-frame extinction. The jet appears to be significantly far from equipartition and magnetically dominated. More extreme versions of this might explain the darkness of many afterglows in the Swift era.

Subject headings: dust, extinction — gamma rays: bursts — magnetic fields — scattering — stars: winds, outflows — stars: Wolf-Rayet

1. Introduction

Discovery of GRB afterglows has become almost commonplace. However, we are still in a regime where nearly every well-sampled afterglow contributes to our understanding of the phenomenon in new and meaningful ways. Observationally, GRB 021211 distinguishes itself in two ways: (1) It is the second GRB for which an optical afterglow was observed within minutes of the burst, thanks to rapid responses by the HETE-2 satellite (Crew et al. 2002, 2003) and three robotic telescopes – RAPTOR (Wozniak et al. 2002); KAIT (Li et al. 2002, 2003); and Super-LOTIS (Park, Williams & Barthelmy 2002; this paper); and (2) It is the first GRB for which filtered detections (beginning 143 seconds after the burst) and color information (beginning 38 minutes after the burst) were obtained at early times.

In addition to observations presented in GRB Coordinates Network (GCN) Circulars, many groups have presented their observations in peer-reviewed journals: Li et al. (2003) present an unfiltered light curve beginning 105 seconds after the burst; Fox et al. (2003) present an unfiltered light curve beginning 21 minutes after the burst and filtered optical, NIR, and radio observations beginning 2.0 hours after the burst; Pandey et al. (2003) present filtered optical observations beginning 6.8 hours after the burst; Holland et al. (2004) present filtered optical and NIR observations of both the afterglow and host galaxy beginning 17 hours after the burst and measure the spectral flux distribution of the afterglow around 21 hours after the burst; and Smith et al. (2005) present submillimeter observations around 25 hours and 10 days after the burst. Finally, Della Valle et al. (2003) present photometric and spectral evidence for an associated supernova at late times.

As in the case of GRB 990123 (Akerlof et al. 1999), the optical afterglow faded more rapidly at first, presumably due to a reverse shock (Wei 2003; Fox et al. 2003; Li et al. 2003; Holland et al. 2004). However, these afterglows differ in that GRB 021211 was $\approx 3 - 4$ mag fainter, despite a lower redshift [$z = 1.004$ for GRB 021211 (Vreeswijk et al. 2002; Della Valle et al. 2002) vs. $z = 1.600$ for GRB 990123 (Hjorth et al. 1999)] (Fox et al. 2003; Li et al. 2003; Pandey et al. 2003; Crew et al. 2003). If it were not for the rapid response of the GRB community, GRB 021211 might have been called a “dark burst”: It faded from $R \approx 14$ mag at ≈ 90 sec after the burst (Wozniak et al. 2002) to $R > 21$ mag about three hours later. Many bursts that would have been called “dark” in the BeppoSAX era are being and will be called “dim” in the HETE-2, Integral, and Swift era due to faster responses.

Some authors have modeled GRB 021211 with an emphasis on its environment. Kumar & Panaitescu (2003) argue that the GRB and afterglow were produced by the same shock and within this framework constrain physical parameters for both constant-density and wind-swept media. Panaitescu & Kumar (2004) consider the early-time afterglows of both GRB 021211 and GRB 990123 in the context of reverse-forward shock (for both constant-density and wind-swept media) and wind-bubble scenarios and find that the reverse-forward shock scenario is preferred. Chevalier, Li & Fransson (2004) argue for a wind-swept medium with the cooling break redward of the R band and within this framework find wind densities that are low compared to Galactic Wolf-Rayet stars (see also Panaitescu & Kumar 2004). Finally, Dado, Dar & De Rujula (2003) model GRB 021211 within the framework of their cannonball model.

In §2, we present FUN GRB Collaboration observations of GRB 021211, which include the earliest filtered detections and color information for an afterglow in the pre-Swift era. In §3, we fit standard afterglow and extinction curve models to these and other

groups’ data and show that within the first few hours after the burst one and possibly two significant chromatic variations are superimposed on this “standard model” behavior. In §4, we compare our results to previous modeling results and discuss possible reasons for these chromatic variations, including the possibility that they are due to a dust echo. We also constrain physical parameters that describe the afterglow and surrounding medium for a variety of scenarios and discuss why GRB 021211’s afterglow is so faint. We draw conclusions in §5.

2. Observations

Long-duration, X-ray rich GRB 021211 was detected by HETE-2’s FREGATE, WXM, and SXC instruments on December 11, 2002 at 11:18:34 UTC (Crew et al. 2003). The initial spacecraft localization was $14'$ in radius and reported in near-real time, only 22 seconds after the burst. Ground analysis of the WXM and SXC data, reported 131 minutes after the burst, improved the localization to $2'$ in radius and was consistent with the initial localization.

Fox & Price (2002) announced the discovery of an $R \sim 18$ mag and fading, stationary point source in the error circle 53 minutes after the burst. While the pair labored, the robotic telescopes of three groups had already responded to the alert. For only the second time in the afterglow era, robotic telescopes extended the light curve of an afterglow back to within tens of seconds of the burst (Wozniak et al. 2002; Li et al. 2002, 2003; Park, Williams & Barthelmy 2002; this paper).

The dim and quickly fading afterglow soon grew too faint for small telescopes, and a possible host galaxy was detected (Lamb et al. 2002a, 2002b; McLeod et al. 2002) but later confirmed under better seeing conditions to be cleanly separated from the afterglow by $1.5''$ (Caldwell et al. 2002). VLT spectroscopy of the true host galaxy resulted in a measured redshift of $z = 1.004 \pm 0.002$ (Vreeswijk et al. 2002; Della Valle et al. 2002). Late-time observations indicate both a re-brightening at the time expected for a supernova at $z \sim 1$, and a spectrum that resembles that of Type Ic SN 1994I (Fruchter et al. 2002; Della Valle et al. 2003).

2.1. FUN GRB Collaboration Observations

We summarize FUN GRB Collaboration observations of GRB 021211 in Table 1. We have calibrated all of our measurements using the field calibration of Henden (2002).

Super-LOTIS imaged the entire GRB 021211 field in R band beginning 143 and 309 seconds after the burst (Park, Williams & Barthelmy 2002). Super-LOTIS is a fully automated f/3.5 0.6-meter diameter Perkin-Elmer telescope on a Boller & Chivens mount at Kitt Peak National Observatory. The camera is a 2048×2048 Loral CCD, which yields a large, $51' \times 51'$ field of view. Observations began automatically after receiving the HETE-2 alert via a socket connection to the GCN. Both exposures were 60 seconds in duration. The mean times that we list in Table 1 are flux weighted using an iterated power-law index of $\alpha = -1.37$, since the exposure time is comparable to the age of the burst, at least for the first exposure. This results in small shifts of 2.4 and 1.2 seconds in the mean times of these observations.

Tenagra Observatories', Ltd., 0.81-meter Tenegra II telescope imaged the GRB 021211 field beginning 37 minutes after the burst. We obtained four sets of four images, each set in a 2×2 arrangement to cover the initial $28'$ -diameter localization and each in a different filter ($I_c R_c V B$). We then re-pointed to the candidate afterglow of Fox & Price (2002) and cycled through $I_c R_c$ thrice more. Of these, we combined the first two I_c and R_c images to optimize signal to noise, but the final two images were not usable due to the onset of morning. This resulted in three detections ($I_c R_c V$), a limit (B), and two more detections ($I_c R_c$). We reduced the images using IRAF's CCDRED package and performed PSF photometry using IRAF's DAOPHOT package.

We imaged the central $11' \times 11'$ of the initial $28'$ -diameter localization in R_c band beginning 85 minutes after the burst from Gunma Astronomical Observatory, located in Agatsuma, Gunma, Japan (Kinugasa et al. 2002). We used the f/12 0.65-meter diameter Cassegrain telescope, which is equipped with an Apogee AP8 1024×1024 back-illuminated SITe CCD. We obtained a total of 28 images, which we combined to optimize signal to noise. We reduced the images using IRAF's CCDRED package and performed PSF photometry on the combined image using IRAF's DAOPHOT package.

We reacquired the field with the 1.34-meter diameter Tautenburg Schmidt telescope 11.7 hours after the burst and imaged in R and I bands for the next 1.1 hours using the 2048×2048 prime-focus CCD (Klose et al. 2002). However, we did not detect the afterglow.

We began observations with the 3.5-meter diameter Astrophysics Research Consortium (ARC) telescope at Apache Point Observatory 22.0 hours after the burst, and returned to the field on December 28 and March 23, 17 and 102 days after the burst (Lamb et al. 2002a, 2002b). All images were taken in i^* band using SPICam, a 2048×2048 back-illuminated SITe CCD. Three 2000-second images were taken on the first night, and seven 1200-second images were taken on each of the following nights. We reduced, combined, and calibrated these images using IRAF's CCDRED and DAOPHOT packages.

Finally, we re-observed the field on December 13 with the 1.0-meter diameter telescope at the U.S. Naval Observatory’s Flagstaff Station for purposes of calibration (Henden 2002). BVR_cI_c images were taken with a 2048×2048 back-illuminated SITe/Tektronix CCD under $2.2''$ seeing conditions. Upon inspection of the images, the afterglow was still marginally visible in the 8-minute V-band image. The afterglow was measured using a two-FWHM diameter aperture with IRAF’s DAOPHOT package.

2.2. Implications of Late-Time ARC Observations

Supernova signatures had been found for many GRBs prior to GRB 021211 (e.g., Galama et al. 1998; Bloom et al. 1999; Reichart 1999; Galama et al. 2000; Bloom et al. 2002; Garnavich et al. 2003; Price et al. 2003; Stanek et al. 2003; see also Zeh, Klose & Hartmann 2004 for a systematic analysis). For GRB 021211, Fruchter et al. (2002) and Della Valle et al. (2003) found evidence for excess red light ≈ 25 days after the burst, and Della Valle et al. (2003) obtained a VLT spectrum at 27 days. This spectrum exhibits Ca II absorption with a relative velocity of $\approx 14,440$ km/s for $z = 1.004$ and is similar to other Type Ic spectra.

Our late-time ARC observations neither confirm nor contradict the existence of this underlying supernova. Subtraction of our second and third i^* epochs using ISIS-2 (Alard 2000) does not reveal any residual flux. However, this is likely due to the timing of our observations: The re-brightening reported by Della Valle et al. (2003) occurs mostly between our observations at 17 and 102 days after the burst. In Figure 1, we plot our i^* light curve and the fitted afterglow model of §3.1.

2.3. Recalibration of KAIT and NEAT Photometry

To better investigate possible chromatic variations that occurred during the unfiltered KAIT and NEAT observations (see §3.2), we have recalibrated these measurements from the R_c band to broad bands given by the spectral responses of their respective CCDs (Pravdo et al. 1999; Li et al. 2003): W. Li (private communication) and P. Price (private communication) kindly provided us with their calibration stars. Using the BVR_cI_c magnitudes of these stars from Henden (2002), we fitted extinguished blackbody functions to each of these stars and then integrated these fitted functions against the appropriate spectral response curve. This resulted in small, 0.05 and 0.03 magnitude offsets in the calibration of the KAIT and NEAT measurements, respectively.

3. Analysis

We now fit standard afterglow and extinction curve models to these and other groups’ data and show that within the first few hours after the burst one and possibly two significant chromatic variations are superimposed on this “standard model” behavior. The data that we include in this analysis are plotted in Figure 2 and consist of FUN GRB Collaboration data (§2.1), data previously published in peer-reviewed journals (Pandey et al. 2003; Fox et al. 2003; Li et al. 2003; Holland et al. 2004), and data from the GCN archive (McLeod et al. 2002). These data span the first ≈ 2.5 days after the burst, after which the host galaxy and supernova become contaminants. All magnitudes have been converted to spectral fluxes as prescribed by Bessell (1979) and Bessell & Brett (1998).

3.1. Model and Fits

We now model these data and constrain model parameters. We model the afterglow with two components, corresponding to reverse and forward shocks. Each component has a power-law light curve and a power-law spectrum, but the spectrum is extinguished by dust in the source frame and in our Galaxy and absorbed by hydrogen in the source frame and the Ly α forest:

$$F_\nu(t) = e^{-\tau_\nu^{MW}} e^{-\tau_{\nu(1+z)}^{Ly\alpha}} e^{-\tau_{\nu(1+z)}^{source}} F_0 \left[\left(\frac{t}{t_0} \right)^{\alpha_{rs}} \left(\frac{\nu}{\nu_R} \right)^{\beta_{rs}} + \left(\frac{t}{t_0} \right)^{\alpha_{fs}} \left(\frac{\nu}{\nu_R} \right)^{\beta_{fs}} \right], \quad (1)$$

where τ_ν^{MW} is the Galactic extinction curve model of Cardelli, Clayton & Mathis (1989), $\tau_{\nu(1+z)}^{Ly\alpha}$ is the Ly α forest absorption model of Reichart (2001a), $\tau_{\nu(1+z)}^{source}$ is the source-frame extinction curve and Lyman limit absorption model of Reichart (2001a), α_{rs} and α_{fs} are the temporal indices of the two components, β_{rs} and β_{fs} are the spectral indices of the two components, ν_R is the effective frequency of the R band, t_0 is the time when these two components are of equal brightness at this frequency, and F_0 is this brightness. Since the extinction and absorption models have features that are narrower than most photometric bands, we integrate Equation 1 against the appropriate filter transmissivity curve (or CCD spectral response curve for the unfiltered measurements; §2.3) before fitting it to the data.

We fit this model to the data using Bayesian inference (e.g., Reichart 2001a; Lee et al. 2001; Galama et al. 2003): The posterior probability distribution is equal to the product of the prior probability distribution and the likelihood function. The likelihood function is given by:

$$\mathcal{L} = \prod_{i=1}^N \frac{1}{\sqrt{2\pi(\sigma_i^2 + \sigma^2)}} \exp \left\{ -\frac{1}{2} \frac{[y(\nu_i, t_i) - y_i]^2}{\sigma_i^2 + \sigma^2} \right\}, \quad (2)$$

where N is the number of measurements, $y(\nu_i, t_i)$ is the above described integration of Equation 1 against the spectral curve of the i th measurement at the time of the i th measurement; y_i is the i th measurement in units of log spectral flux; σ_i is the uncertainty in the i th measurement in the same units, and σ is a parameter, sometimes called the slop parameter, that models the small systematic errors that are unavoidably introduced when data are collected from many sources, and other small sources of error (Reichart 2001a). Ignoring this parameter can lead to erroneous fits and significantly underestimated uncertainties in the fitted parameter values when the scatter of the measurements about the fitted model exceeds that which can be accounted for by the measurement uncertainties alone.

Many of the parameters of the source-frame extinction curve model and all of the parameters of the Ly α forest absorption and Galactic extinction curve models can be constrained a priori. The source-frame extinction curve model of Reichart (2001a) is a function of eight parameters: the source-frame V-band extinction magnitude A_V , $R_V = A_V/E(B - V)$, the intercept c_1 and slope c_2 of the linear component of the source-frame UV extinction curve, the strength c_3 , width γ and center x_0 of the UV bump component of the extinction curve, and the strength c_4 of the FUV excess component of the extinction curve. The Ly α forest absorption model of Reichart (2001a) is a function of a single parameter, D_A , the flux deficit. Reichart (2001a) determines prior probability distributions for R_V , c_1 , γ , x_0 , and D_A , which means that the values of these parameters can be weighted by fairly narrow distributions, the description of which sometimes depends on other parameters (c_2 and z), a priori. We adopt these priors here, which can be thought of as increasing the degrees of freedom by five. Also, the Galactic extinction curve model of Cardelli, Clayton & Mathis (1989) is a function of $E(B - V) = 0.028$ mag for this line of sight (Schlegel, Finkbeiner & Davis 1998) and a single parameter, R_V^{MW} . We adopt a prior for this parameter that is log normally distributed with mean log 3.1 and width 0.1, which closely approximates the distribution of values of this parameter along random lines of sight through the Galaxy (e.g., Reichart 2001a; Lee et al. 2002; Galama et al. 2003).

We fit our model to the data for each of the four standard cases of Sari, Piran & Narayan (1998) and Chevalier & Li (2000), which relate α_{fs} to β_{fs} assuming (1) propagation through either a constant-density (ISM) or wind-swept (WIND) medium, and (2) a cooling break that is either redward (RED) or blueward (BLUE) of the observed optical and NIR frequencies: For the ISM-RED and WIND-RED cases, $\alpha_{fs} = (3\beta_{fs} + 1)/2 = -(3p - 2)/4$; for the ISM-BLUE case, $\alpha_{fs} = 3\beta_{fs}/2 = -3(p - 1)/4$; and for the WIND-BLUE case, $\alpha_{fs} = (3\beta_{fs} - 1)/2 = -(3p - 1)/4$, where p is the power-law index of the electron-energy distribution. Since the temporal index is well constrained by the data, these additional constraints can be powerful tools for separating the intrinsic spectrum from extinction

effects (see §4.1). For purposes of comparison, we also fit our model to the data free of constraints on α_{fs} and β_{fs} .

Best fits are found by maximizing the posterior. Compared to the WIND-BLUE case, we can rule out the ISM-RED and WIND-RED cases at the 7.3σ credible level, and the ISM-BLUE case is disfavored at the 3.1σ credible level. Furthermore, the WIND-BLUE fit is consistent with the constraint-free fit, differing from it at only the 0.6σ credible level. The primary difference between these cases is that the WIND-BLUE case requests a shallow intrinsic spectrum, $\beta_{fs} = -0.34^{+0.01}_{-0.01}$, and a small amount of extinction, $A_V = 0.18^{+0.25}_{-0.12}$ mag, where the other cases request steeper intrinsic spectra and would fit better if $A_V < 0$ mag were possible (see §4). For the WIND-BLUE case, we find that $\log F_0 = 2.98^{+0.12}_{-0.12}$ μJy , $\log t_0 = -2.56^{+0.07}_{-0.07}$ day, $\alpha_{rs} = -2.16^{+0.09}_{-0.10}$, $\beta_{rs} = 1.1^{+0.7}_{-0.8}$, $\alpha_{fs} = -1.01^{+0.02}_{-0.01}$, $\beta_{fs} = -0.34^{+0.01}_{-0.01}$, $A_V = 0.18^{+0.25}_{-0.12}$ mag, $c_2 < 4.3$ (1σ), and $\sigma = 0.038^{+0.010}_{-0.008}$ mag.¹⁹ The parameters c_3 and c_4 could not be constrained by the data. We plot best-fit light curves for 13 spectral bands in Figure 2 and best-fit spectral flux distributions for six epochs in Figure 4.

3.2. Chromatic Variations

We plot the residuals of Figure 2 in Figure 3. One and possibly two significant chromatic variations can be seen from ≈ 40 minutes after the burst until possibly ≈ 6.0 hours after the burst. The first of these is an increase relative to the best-fit model of the unfiltered NEAT and KAIT data, which is also clearly visible in Figure 2b, concurrent with a decrease relative to the best-fit model of our R_c and possibly I_c data from Tenagra and Gunma. Since the NEAT and KAIT bandpasses are broad, encompassing the I_c and R_c bands on their red ends, this suggests that there was an excess of blue light at this time. To explore this further, we plot the spectral flux distribution of the afterglow in six time slices in Figure 4:

In Figure 4a, we plot the best-fit spectral flux distribution at 67 minutes after the burst and have scaled all of the data between 39 and 94 minutes after the burst to this time using the best-fit light curve. These data consist of $I_c R_c$ VB data from Tenagra and unfiltered NEAT data. We plot the weighted average of the scaled NEAT data for clarity. The combined NEAT point is only 0.14 mag above the best-fit model, but significantly so, at the 5.2σ confidence level.

¹⁹Due to the dimension of the parameter space, marginalized probability distributions for each parameter value would take impossibly long to compute. Consequently, these error bars are measured from projected probability distributions and are consequently conservative overestimates.

In Figure 4b, we plot the best-fit spectral flux distribution at 2.2 hours after the burst. We have scaled all of the data between 1.8 and 2.5 hours after the burst to this time using the best-fit light curve and plot weighted averages of the scaled data when there are multiple points per spectral band. These data consist of a K_s point from Fox et al. (2003), $I_c R_c$ data from Tenagra, an R_c point from Gunma, and unfiltered KAIT data. The combined KAIT point is even farther above the best-fit model, 0.35 mag, this time at the 3.3σ confidence level. The I_c point is now below but still consistent with the best-fit model and the R_c point is below the best-fit model at the 2.3σ confidence level. Consequently, the KAIT point differs from the R_c point at the 4.0σ confidence level with respect to the best-fit model. Since the KAIT bandpass, like the NEAT bandpass, is broad, encompassing the I_c and R_c bands on its red end, this suggests that there was an excess of blue light at this time. If we model this excess as an additional power-law component, just in this time slice, we find it to be bluer than $\beta = 1.0$ at the 2σ credible level.

In Figure 4c, we plot the best-fit spectral flux distribution at 4.4 hours after the burst. We have scaled two points from Fox et al. (2003) – a B point at 3.1 hours after the burst and an R_c point at 5.7 hours after the burst – to this time using the best-fit light curve. One possibility is that the excess light has changed from blue to red: The R_c point is above the best-fit model at the 3.1σ confidence level and the B point is below the best-fit model at the 3.7σ confidence level. Consequently, these points differ at the 4.8σ confidence level with respect to the best-fit model. However, given the sparsity of the data in this time slice a temporal variation cannot be ruled out either.

The remaining panels, corresponding to 6.8 – 11, 17 – 25, and 46 – 48 hours after the burst, show no evidence for significant chromatic variations at later times.

Although the third time slice is too sparsely sampled for a temporal variation to be ruled out, the first two time slices, which span the first proposed chromatic variation, are better sampled. Consider the following simple model: Let t_1 be the beginning of this variation. Prior to t_1 , the afterglow is described by Equation 1. Between t_1 and 2.5 hours after the burst, the NEAT and KAIT data are instead described by temporal index α_{NK} and the R_c and I_c data are instead described by temporal index $\alpha_{R_c I_c}$. If this were a temporal variation, α_{NK} would equal $\alpha_{R_c I_c}$. Instead, we find that $t_1 = 46_{-21}^{+14}$ min and $\alpha_{NK} - \alpha_{R_c I_c} = 0.46_{-0.19}^{+0.23}$ with $\alpha_{NK} - \alpha_{R_c I_c} > 0$ at the 3.5σ credible level. Here, we have fixed all of the other parameters to their previous best-fit values so we can also plot this best fit in Figure 3. Allowing all of the parameters to vary, we find that $\alpha_{NK} - \alpha_{R_c I_c} > 0$ at the 3.3σ credible level, which again suggests that this is a chromatic variation.

In the above fit, we also find that $\alpha_{R_c I_c} = -1.30_{-0.26}^{+0.20}$, which is somewhat steeper than the fitted value of α_{fs} . This suggests that the light curve might be steepening during

the first and second time slices, and this is consistent with the B point at the beginning of the third time slice also undercutting the model (e.g., Figure 4c). However, the data are consistent with the model in the fourth, fifth, and sixth time slices, which suggests a minor rebrightening during the third time slice. Such minor temporal variations are now commonplace – GRBs 021004 and 030329 are extreme examples – but further modeling of such variations is beyond the scope of this paper, and frankly beyond the quality of this data set. However, this does lend some credibility to the possibility that the second variation is temporal instead of chromatic. A final possibility is that the R point at the end of the third time slice is a statistical variation: Given 80 points, the probability of encountering a 4.8σ variation is 1 in 7900 (ruled out at the 3.8σ confidence level).

Finally, we refit the four standard cases to the data, but this time we accommodate the first, chromatic variation with the above simple model and eliminate the second variation, whether chromatic or temporal, by not fitting to the two points of the third time slice. Compared to the WIND-BLUE case, we now rule out the ISM-RED and WIND-RED cases at the 6.3σ credible level and the ISM-BLUE case at the 3.2σ credible level. For the WIND-BLUE case, we find that $\log F_0 = 2.27^{+0.26}_{-0.25} \mu\text{Jy}$, $\log t_0 = -2.01^{+0.17}_{-0.16}$ day, $\alpha_{rs} = -1.78^{+0.07}_{-0.08}$, $\beta_{rs} = -0.95^{+1.45}_{-0.55}$, $\alpha_{fs} = -0.89^{+0.04}_{-0.04}$, $\beta_{fs} = -0.26^{+0.03}_{-0.03}$, $A_V = 0.35^{+0.22}_{-0.17}$ mag, $c_2 < 1.6$ (1σ), $t_1 = 7.5^{+2.7}_{-3.5}$ min, $\alpha_{RcIc} = -0.88^{+0.08}_{-0.08}$, $\alpha_{NK} - \alpha_{RcIc} = 0.12^{+0.06}_{-0.06}$, and $\sigma = 0.028^{+0.007}_{-0.007}$ mag. The primary difference between this fit and the WIND-BLUE fit of §3.1 in which these variations are not treated is we now find more source-frame extinction, $A_V = 0.35^{+0.22}_{-0.17}$ mag with $A_V > 0$ mag at the 2.8σ credible level. Also, α_{RcIc} is now consistent with α_{fs} , but $\alpha_{NK} - \alpha_{RcIc}$ is still greater than zero at the 3.0σ credible level.

4. Discussion

4.1. Model and Fits

Our finding that the data are best described by the WIND-BLUE case differs from the findings of others. Fox et al. (2003) discount this case in favor of the ISM-BLUE case, arguing that if the early-time emission is due to a reverse shock, in a wind-swept medium it is expected to fade quickly and they measure a slower fading: $\alpha_{rs} = -1.63 \pm 0.13$. However, Chevalier, Li & Fransson (2004) point out that this measurement depends sensitively on how one subtracts out (or models) the forward-shock component, arguing that the value is closer to $\alpha_{rs} = -2.2$. Using final instead of GCN data, we find that $\alpha_{rs} = -2.16^{+0.09}_{-0.10}$ (variations untreated; §3.1) or $-1.78^{+0.07}_{-0.08}$ (variations treated; §3.2). However, in §4.2 we point out that emission from the reverse shock is not necessarily expected to fade quickly in a wind-swept medium if A_* and other physical parameters are lower than expected, which

appears to be the case for this GRB.

Holland et al. (2004) also adopt the ISM-BLUE case. The primary difference between their fit and ours is that we permit source-frame extinction. When we fit the ISM-BLUE case, we find that $\beta_{fs} = -0.67$ (variations untreated) or -0.60 (variations treated) with $A_V = 0$ mag, which is very similar to their fit in a time slice around 0.88 days after the burst: $\beta_{fs} = -0.69 \pm 0.14$ with A_V assumed to be zero. However, if source-frame extinction is permitted and one fits to all of the data, we find that the WIND-BLUE case with a small to moderate amount of source-frame extinction, $A_V = 0.18^{+0.25}_{-0.12}$ mag (variations untreated) or $0.35^{+0.22}_{-0.17}$ mag (variations treated), is preferred at the 3.1σ (variations untreated) or 3.2σ (variations treated) credible level. Figures 4e and 5e can be directly compared to Figure 3 of Holland et al. (2004).

Finally, Chevalier, Li & Fransson (2004) adopt the WIND-RED case, guided by sparse color information that was available at the time, including the two points of Figure 4c, which we have already identified as discrepant, possibly due to excess red light at this time (§3.2). Permitting source-frame extinction and fitting to all of the data, we rule this case out at the 7.3σ (variations untreated) or 6.3σ (variations treated) credible level.

The WIND-BLUE case, however, requires a relatively hard electron energy distribution $-p = 1.68^{+0.01}_{-0.03}$ – so a break at higher energies is required. Bhattacharya (2001) determines the effect of $p < 2$ on the standard equations: By introducing a cut-off frequency γ_u such that $\gamma_m < \gamma_e < \gamma_u$ and assuming that γ_u evolves directly with the bulk Lorentz factor of the shock, they find results similar to the standard prescriptions. Galama et al. (2003) found a similar hard electron energy index for GRB 010222, though other ideas, such as a continuous injection of energy (Bjornsson et al. 2002) or an early transition to non-relativistic motion (in’t Zand et al. 2001; Masetti et al. 2001), have been proposed.

4.2. Physical Parameters

Following the analysis of Chevalier, Li & Fransson (2004), but for the WIND-BLUE case, and using the analytic expressions of Granot & Sari (2002), we now constrain physical parameters that describe the afterglow and surrounding medium for a variety of scenarios. The first constraint comes from the expression of Granot & Sari (2002) for the brightness of the afterglow in the frequency range of our observations, which for the WIND-BLUE case is $\max\{\nu_{sa}, \nu_m\} < \nu < \nu_c$, where ν_{sa} is the self-absorption frequency, ν_m is the typical synchrotron frequency, and ν_c is the electron cooling frequency. This corresponds to segment G in their Figure 1. For $p = 1.68$, a luminosity distance of $d_L = 2.06 \times 10^{28}$ cm (assuming

that $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$), and an extinction-corrected $F_R = 19 \text{ } \mu\text{Jy}$ at 0.1 days after the burst, we find:

$$\bar{\epsilon}_e^{0.68} \epsilon_B^{0.67} A_* E_{52}^{0.67} = 1.02 \times 10^{-5}, \quad (3)$$

where $\bar{\epsilon}_e$ is the electron energy fraction when $p < 2$, ϵ_B is the magnetic field energy fraction, A_* measures the density of the wind-swept medium, and $E = E_{52} \times 10^{52} \text{ erg}$ is the total energy of the shock, if spherical. The second constraint comes from the expression of Granot & Sari (2002) for $\nu_c(t)$ and the fact that the data are well described by the WIND-BLUE case even at early times (see §4.3). Taking $\nu_c > \nu_R$ prior to 3.9 minutes after the burst – the time when the forward shock first outshines the reverse shock in the R band – yields:

$$\epsilon_B^{3/2} A_*^2 E_{52}^{-1/2} = 6.52 \times 10^{-6} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{1/2}, \quad (4)$$

where $t_{c,R}$ is the time that ν_c passes above the R band. The third constraint is similar to the second in that we take $\nu_m < \nu_R$ prior to 3.9 minutes after the burst, else the light curve would have faded much more slowly at this time, as $F_\nu \sim t^{-1/4}$ (Chevalier & Li 2000; Chevalier, Li & Fransson 2004):

$$E_{52}^{1/2} \bar{\epsilon}_e^2 \epsilon_B^{1/2} = 1.18 \times 10^{-5} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{3/2}, \quad (5)$$

where $t_{m,R}$ is the time that ν_m passes below the R band. The final constraint comes from the expression of Granot & Sari (2002) for the brightness of the afterglow at 8.5 GHz, given that $F_{8.5} < 35 \text{ } \mu\text{Jy}$ at a mean time of 13 days after the burst (Fox et al. 2003). Here we consider four scenarios: (A) $\nu_{sa} < 8.5 \text{ GHz} < \nu_m$, (B) $8.5 \text{ GHz} < \min\{\nu_{sa}, \nu_m\}$, (C) $\max\{\nu_{sa}, \nu_m\} < 8.5 \text{ GHz}$, and (D) $\nu_m < 8.5 \text{ GHz} < \nu_{sa}$.

For scenario A, using the expression of Granot & Sari (2002) that corresponds to their segment D, we find:

$$\bar{\epsilon}_e^{-2/3} \epsilon_B^{1/3} A_* E_{52}^{1/3} = 3.76 \times 10^{-2} \left(\frac{F_{8.5}}{35 \text{ } \mu\text{Jy}} \right). \quad (6)$$

Combining Equations 3, 4, 5, and 6 yields:

$$\bar{\epsilon}_e = 6.53 \times 10^{-4} E_{52}^{-1} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.25} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{0.24}, \quad (7)$$

$$\epsilon_B = 765 E_{52}^3 \left(\frac{t_{c,R}}{3.9 \text{ min}} \right) \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{2.04}, \quad (8)$$

$$A_* = 1.75 \times 10^{-5} E_{52}^{-2} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.5} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-1.53}, \quad (9)$$

$$F_{8.5} = 20 \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-1.01} \mu\text{Jy}. \quad (10)$$

If we additionally require that $\nu_{sa} < 8.5 \text{ GHz} < \nu_m$ at 13 days after the burst, using the expressions of Granot & Sari (2002) for ν_{sa} and ν_m , we find that $F_{8.5} < 6 \mu\text{Jy}$ and $E_{52} > 3.50 \times 10^{-3} (F_{8.5}/35 \mu\text{Jy})^{1.38}$. Since the former of these constraints contradicts Equation 10 for any value of $t_{m,R} < 3.9 \text{ min}$, we rule out this scenario.

For scenario B, using the expression of Granot & Sari (2002) that corresponds to their segment B, we find:

$$\bar{\epsilon}_e A_*^{-1} E_{52} = 1.70 \times 10^{-4} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right). \quad (11)$$

Combining Equations 3, 4, 5, and 11 yields:

$$\bar{\epsilon}_e = 0.306 \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.125} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{1.125} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{-0.5}, \quad (12)$$

$$\epsilon_B = 7.45 \times 10^{-6} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{0.625} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-0.615} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{1.5}, \quad (13)$$

$$A_* = 3.85 \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.25} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{0.24} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{-1}, \quad (14)$$

$$E_{52} = 2.14 \times 10^{-3} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.125} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-0.885} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{0.5}. \quad (15)$$

If we additionally require that $8.5 \text{ GHz} < \min\{\nu_{sa}, \nu_m\}$ at 13 days after the burst, we find that $t_{m,R} > 2.9 \text{ min}$, which is technically consistent with $t_{m,R} < 3.9 \text{ min}$, and $F_{8.5} > 27 \mu\text{Jy}$, which is technically consistent with $F_{8.5} < 35 \mu\text{Jy}$. However, this constrains these parameters' values to narrow ranges and by Equation 15 implies a value for E_{52} that is much too low, given that the isotropic-equivalent energy in gamma rays alone was $(1.0 \pm 0.1) \times 10^{52} \text{ erg}$ (Holland et al. 2004) or $1.68_{-0.27}^{+0.32} \times 10^{52} \text{ erg}$ (Lamb et al. 2004). Assuming that the efficiency at which energy is converted to gamma rays is $\sim 20\%$ (e.g., Beloborodov 2000), then $E_{52} \sim \text{many}$. Consequently, we rule out this scenario as well.

For scenario C, using the expression of Granot & Sari (2002) that corresponds to their segment G, we find:

$$\bar{\epsilon}_e^{0.68} \epsilon_B^{0.67} A_* E_{52}^{0.67} = 6.26 \times 10^{-5} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right). \quad (16)$$

Combining Equations 3, 4, 5 and 16 yields:

$$\bar{\epsilon}_e = 6.53 \times 10^{-4} E_{52}^{-1} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.25} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{0.24}, \quad (17)$$

$$\epsilon_B = 765 E_{52}^3 \left(\frac{t_{c,R}}{3.9 \text{ min}} \right) \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{2.04}, \quad (18)$$

$$A_* = 1.75 \times 10^{-5} E_{52}^{-2} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.5} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-1.53}, \quad (19)$$

$$F_{8.5} = 6 \mu\text{Jy}. \quad (20)$$

If we additionally require that $\max\{\nu_{sa}, \nu_m\} < 8.5 \text{ GHz}$ at 13 days after the burst, we find that $t_{m,R} < 13 \text{ minutes}$ and $E_{52} > 1.27 \times 10^{-3}$, neither of which are problematic. Taking E_{52} to be as low as 4 and $t_{c,R}$ and $t_{m,R}$ to be as low as the duration of the burst ($T_{90} = 2.41 \pm 0.15 \text{ sec}$ in the 30 – 85 keV band, in which νF_ν peaks; Crew et al. 2003), yields $\epsilon_B \sim 0.04$. In this case, $\epsilon_e \sim 0.0002$ and $A_* \sim 0.01$. However, for ϵ_B to be this low requires considerable fine tuning: If E_{52} is as high as 11, $t_{c,R}$ is as high as 54 sec, $t_{m,R}$ is as high as 11 sec, or lesser combinations of these three, $\epsilon_B \approx 1$. Consequently, ϵ_B is likely considerably more than 0.04, in which case ϵ_e can be no more than a factor of three greater and is likely less and A_* can only be less.

Finally, for scenario D, using the expression of Granot & Sari (2002) that corresponds to their segment A, we find:

$$\epsilon_B^{-1/4} A_*^{-1} E_{52}^{3/4} = 8.60 \times 10^{-2} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right). \quad (21)$$

Combining Equations 3, 4, 5 and 21 yields:

$$\bar{\epsilon}_e = 0.232 \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.125} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{0.75} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{-0.5}, \quad (22)$$

$$\epsilon_B = 1.71 \times 10^{-5} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{0.625} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{0.51} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{1.5}, \quad (23)$$

$$A_* = 2.21 \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.25} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-0.51} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{-1}, \quad (24)$$

$$E_{52} = 2.82 \times 10^{-3} \left(\frac{t_{c,R}}{3.9 \text{ min}} \right)^{-0.125} \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{-0.51} \left(\frac{F_{8.5}}{35 \mu\text{Jy}} \right)^{0.5}, \quad (25)$$

If we additionally require that $\nu_m < 8.5 \text{ GHz} < \nu_{sa}$ at 13 days after the burst, we find that $t_{m,R} > 2.9 \text{ min}$, which is technically consistent with $t_{m,R} < 3.9 \text{ min}$, and $F_{8.5} > 7 \mu\text{Jy}$, which is consistent with $F_{8.5} < 35 \mu\text{Jy}$. Once again, this constrains these parameters' values to relatively narrow ranges and by Equation 25 implies a value for E_{52} that is much too low. Consequently, we rule out this scenario as well.

Consequently, we find that $\nu_m < \nu_R < \nu_c$ after $< 3.9 \text{ minutes}$ after the burst and $\max\{\nu_{sa}, \nu_m\} < 8.5 \text{ GHz}$ around 13 days after the burst. In this scenario, $\bar{\epsilon}_e$ and A_* are

considerably lower than canonical values. Since $F_R \propto \bar{\epsilon}_e^{0.68} \epsilon_B^{0.67} A_* E_{52}^{0.67}$ (Equation 3), both of these contribute to the faintness of the afterglow (§1).

If we allow ourselves to be guided by the standard-energy result (Frail et al. 2001; Bloom, Frail & Kulkarni 2003), E_{52} is also lower than what one might expect: For wind-swept media, the total energy released in gamma rays is typically measured to be many $\times 10^{50}$ erg (Bloom, Frail & Kulkarni 2003). Given that the isotropic-equivalent energy in gamma rays for GRB 021211 was $(1.0 \pm 0.1) \times 10^{52}$ erg (Holland et al. 2004) or $1.68_{-0.27}^{+0.32} \times 10^{52}$ erg (Lamb et al. 2004), this implies a jet opening/viewing angle of $\sim 20^\circ$, which is about three times the canonical value. Hence, E_{52} is probably about an order of magnitude less than what one might have expected. Consequently, we find that GRB 021211’s afterglow is faint for a combination of 3 – 4 reasons: (1) a low fraction of energy in relativistic electrons, (2) a low density for the wind-swept medium, implying either a low mass-loss rate and/or a high wind velocity for the progenitor, (3) a wide opening/viewing angle for the jet, and possibly (4) moderate source-frame extinction (§3.2).

Furthermore, with $\epsilon_B/\bar{\epsilon}_e > 200$ and ϵ_B likely much greater than 0.04 the jet appears to be significantly far from equipartition and magnetically dominated. This is similar to SN 1993J, for which the magnetic field energy density is $\sim 10^4$ times the relativistic particle energy density (Chandra et al. 2004), but dissimilar to SN 1998bw (Kulkarni et al. 1998) and SN 2002ap (Bjornsson & Fransson 2004), which appear to be near equipartition.

These findings are supported by the existence of the bright reverse shock: Fox et al. (2003) dismiss the possibility of a wind-swept medium because for canonical values of $\bar{\epsilon}_e^{rs}$, ϵ_B^{rs} , A_* , and E_{52} , ν_c^{rs} is expected to be significantly less than ν_m^{rs} , in which case the reverse shock is expected to fade away quickly and not be bright (Chevalier & Li 2000). However, this is not the case when $\bar{\epsilon}_e^{rs}$, ϵ_B^{rs} , and/or A_* are sufficiently low. Taking $\bar{\epsilon}_e^{rs} \sim \bar{\epsilon}_e$ and $\epsilon_B^{rs} \sim \epsilon_B$ and substituting Equations 17, 18, and 19 into Equations 45 and 47 of Chevalier & Li (2000), we find:

$$\nu_c^{rs}/\nu_m^{rs} = 5.56 \times 10^4 E_{52}^3 \left(\frac{t_{m,R}}{3.9 \text{ min}} \right)^{0.03} \left(\frac{t}{1 \text{ min}} \right)^2, \quad (26)$$

for $X = 0.75$, $\Delta_{10} = 3$, and $\gamma_3 = 0.3$. For $t = 1$ minute after the burst, corresponding to the beginning of the first detection of the reverse shock (Wozniak et al. 2002), E_{52} need only be greater than ~ 0.03 for $\nu_c^{rs} \sim \nu_m^{rs}$.

4.3. Chromatic Variations

Globally, the data are well described by the model of §3.1, but superimposed on this global behavior are small, but significant variations from ≈ 40 minutes after the burst until possibly ≈ 6.0 hours after the burst (§3.2). This model does not explain these variations. It merely attempts to accommodate them with a higher value of the slop parameter: $\sigma = 0.038^{+0.010}_{-0.008}$ mag. The slop parameter is a global measure of the scatter of the data around the model, beyond what can be accounted for by the data’s error bars (§3.1). Furthermore, at least the first of these variations appears to be chromatic, with a relative increase of blue light with respect to red light around two hours after the burst, and possibly, but less certainly, a reversal of this two hours later (§3.2). When we modify the model and fit to better treat these variations, σ decreases to $0.028^{+0.007}_{-0.007}$ mag (§3.2).

One potential explanation for these variations is that we have undersampled a light curve that is varying in time in such a way as to mimic a chromatic effect. Indeed, the high signal-to-noise light curves of GRBs 021004 and 030329 revealed temporal variations and a variety of explanations have been proposed, including variations in the density of the external medium (e.g., Lazzati et al. 2002), refreshed shocks (e.g., Granot, Nakar & Piran 2003), and patchy shells (e.g., Kumar & Piran 2000). However, none of these occur on a sufficiently short timescale to explain the variation around two hours after the burst (Figure 3). However, a temporal variation cannot be ruled out for the possible reversal of this effect two hours later (§3.2).

Another potential explanation is that a spectral break, presumably the cooling break, is passing through our data around 2 – 4 hours after the burst. Although this is difficult to reconcile with the blue excess of Figure 4b, since the spectrum is supposed to be half of a spectral index steeper blueward of the cooling break, it is not necessarily inconsistent with the possible red excess of Figure 4c. However, the spectra in Figures 4a and 4e would then differ by half of a spectral index, which would be noticeable.

Another potential explanation is that we observed a dust echo – light scattered by dust into the line of sight and received with a time delay due to the greater path length. Waxman & Draine (2000) and Esin & Blandford (2000) originally proposed dust echoes as an alternative explanation for the supernova-like components to the afterglows of GRB 980326 (Bloom et al. 1999) and GRB 970228 (Reichart 1999; Galama et al. 2000). Reichart (2001b) modeled and computed dust echo light curves and spectral flux distributions and found that while dust echoes can mimic supernova light curves they cannot mimic supernova spectral flux distributions, at least not near the spectral peak. Moran & Reichart (2004) take the model of Reichart (2001b) and instead of applying it to dust shells of inner radius $\sim 10^{18}$ cm, which is what is required to mimic supernova light curves, they apply it

to dust shells of inner radius $\sim 10^{14} - 10^{15}$ cm, which is typical of late-type WC Wolf-Rayet stars, the likely progenitors of GRBs.

Moran & Reichart (2004) find that for (1) wind-swept media, (2) bright optical flashes – reverse shocks that outshine the forward shock at early times – and (3) wide jet opening angles, dust echoes may be observable on a timescale of minutes to hours after the burst. Furthermore, the characteristic signature will be an excess of blue light (since blue light scatters preferentially) that quickly transitions to an excess of red light (due to increasing path lengths through dust with increasing time delay). Since all of these conditions appear to be met for GRB 021211 (§3.1, §4.1, §4.2), and excess blue light is observed on this timescale, as well as a possible transition to excess red light hours later, we now test this hypothesis in two ways:

1. Equation 6 of Moran & Reichart (2004) gives the turn-on time of an idealized, on-axis dust echo as a function of the inner radius R of the circum-progenitor dust shell and the initial opening angle θ_{jet} of the jet. Dust echoes should not be visible at very early times because X rays from the burst and UV light from the optical flash should sublimate the dust within θ_{jet} of the jet axis this close to the burst (e.g., Waxman & Draine 2000; Fruchter, Krolik & Rhoads 2001; Reichart 2001c). Taking the turn-on time to be ~ 0.4 of the peak time (Reichart 2001b; Moran & Reichart 2004) and taking the peak time for GRB 021211 to be ~ 2.2 hours, we solve for R :

$$R \sim 3 \times 10^{15} \left(\frac{\theta_{jet}}{10^\circ} \right)^{-2} \text{ cm.} \quad (27)$$

For a wide jet opening angle, this yields $R \sim 10^{14} - 10^{15}$ cm, which is the expectation if late-type WC stars are indeed the progenitors of GRBs. However, given that the isotropic-equivalent energy in gamma rays alone was $(1.0 \pm 0.1) \times 10^{52}$ erg (Holland et al. 2004) or $1.68_{-0.27}^{+0.32} \times 10^{52}$ erg (Lamb et al. 2004) and that $A_* \lesssim 0.01$ (§4.2), the deceleration radius r_d was likely greater than 10^{16} cm (e.g., Equation 7 of Moran & Reichart 2005) and consequently a value for R likely cannot be deduced.

2. Equation 5 of Moran & Reichart (2004), but with r_d substituted for R , gives the peak brightness of the optical flash off of the jet axis, at angles around θ_{jet} , as a function of the peak brightness of the dust echo, the deceleration radius, the optical depth $\tau_{\nu(1+z)}$ through the rest of the dust shell at frequency $\nu(1+z)$, also at angles around θ_{jet} , and the duration of the optical flash. Taking the peak brightness of the dust echo to be ~ 19 mag and the duration of the optical flash to be ~ 30 seconds (e.g., Sari & Piran 1999), we find:

$$m_{OF}(\theta \sim \theta_{jet}) \sim 9 + 2.5 \log \tau_{\nu(1+z)} - 2.5 \log \left(\frac{r_d}{10^{16} \text{ cm}} \right) \text{ mag,} \quad (28)$$

Using our best fit (§3.1), extrapolation of the reverse shock light curve back to ≈ 30 seconds after the onset of the burst yields $R \sim 12$ mag. However, given the known distance to GRB 021211 $m_{OF}(\theta \sim \theta_{jet})$ would have to be significantly fainter lest the dust be sublimated in these directions as well, this close to the burst. For longer optical flash durations, $\tau_{\nu(1+z)} > 1$, and/or a greater forward scattering probability than what Esin & Blandford (2000), Reichart (2001b), and Moran & Reichart (2004) assume, $m_{OF}(\theta \sim \theta_{jet})$ would be fainter, but not sufficiently. Alternatively, prior fragmentation of the dust to PAH levels by gamma rays from the burst might harden it against sublimation, since atomic bonds would then be more difficult to break.

Finally, we point out that neither of these estimates hold in the case of a jet with a narrow opening angle but a large viewing angle.

5. Conclusions

GRB 021211 is one of only a handful of GRBs for which processes other than the forward shock have been identified at optical wavelengths, which has made it one of the most studied GRBs. In this paper, we present additional, multi-band observations of this event, ranging from minutes to months after the burst, which in combination with all previously published observations have allowed us to deeply probe the physics of this GRB and properties of its circum-progenitor environment.

Coupling the standard afterglow model with a general-purpose extinction curve model, we find that the afterglow is best described by propagation through a wind-swept medium, which implies a massive-star progenitor (e.g., Price et al. 2002). The jet itself appears to be significantly far from equipartition and magnetically dominated. Indeed, the low fraction of energy in relativistic electrons appears to be the primary reason that this afterglow is so faint. This, combined with a low-density medium, a wide jet opening/viewing angle, and possibly moderate extinction might be important clues as to why many afterglows are dark/dim in the Swift era, even at early times after the burst. These findings are supported by the existence of the bright reverse shock – in a wind-swept medium this should only be possible if A_* is low and/or the jet is significantly far from equipartition, meaning that either $\bar{\epsilon}_e^{rs}$ or ϵ_B^{rs} is low as well.

Finally, we observed one and possibly two significant chromatic variations hours after the burst. We discuss possible reasons for these variations, including the possibility that they are due to a dust echo: The three primary requirements for an observable dust echo are a wind-swept medium, a bright optical flash, and a wide jet opening angle, and

the characteristic signature should be an excess of blue light that quickly reddens, all of which appear to be satisfied for GRB 021211. However, in the case of GRB 021211 this would imply an off-axis brightness and hence luminosity for the optical flash that would probably sublimate too much dust in these directions. Rapid, multi-band, and preferably simultaneous multi-band observations of future GRBs might shed more light on this interesting possibility.

DER very gratefully acknowledges support from NSF's MRI, CAREER, PREST, and REU programs, NASA's APRA, Swift GI and IDEAS programs, UNC's Junior Faculty Development Award, and especially Leonard Goodman and Henry Cox. HSP acknowledges support from NASA's SR&T program and DOE/UC LLNL contract W-7405-Eng-48. KK acknowledges support from Ministry of Education, Culture, Sports, Science and Technology of Japan grant-in-aid 16740121. DQL acknowledges support from NASA subcontract NAGW-4690 (HETE-2) and NASA's LTSA program.

REFERENCES

- Akerlof, C., et al. 1999, *Nature*, 398, 400
- Alard, C. 2000, *A&AS*, 144, 363
- Beloborodov, A. M. 2000, *ApJ*, 539, L25
- Bessell, M. S. 1979, *PASP*, 91, 589
- Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
- Bhattacharya, D. 2001, *Bulletin of the Astronomical Society of India*, 29, 107
- Björnsson, G., Hjorth, J., Pedersen, K., & Fynbo, J. U. 2002, *ApJ*, 579, L59
- Björnsson, C., & Fransson, C. 2004, *ApJ*, 605, 823
- Bloom, J. S., et al. 1999, *Nature*, 401, 453
- Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, *ApJ*, 594, 674
- Caldwell, N., McLeod, B., Garnavich, P., & Stanek, K. Z. 2002, *GRB Circular Network*, 1759, 1
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245

- Chandra, P., Ray, A., & Bhatnagar, S. 2004, *ApJ*, 612, 974
- Chevalier, R. A., & Li, Z. 2000, *ApJ*, 536, 195
- Chevalier, R. A., Li, Z., & Fransson, C. 2004, *ApJ*, 606, 369
- Crew, G., et al. 2002, *GRB Circular Network*, 1734, 1
- Crew, G. B., et al. 2003, *ApJ*, 599, 387
- Dado, S., Dar, A., & De Rújula, A. 2003, *ApJ*, 593, 961
- Della Valle, M., et al. 2002, *GRB Circular Network*, 1809, 1
- Della Valle, M., et al. 2003, *A&A*, 406, L33
- Esin, A. A., & Blandford, R. 2000, *ApJ*, 534, L151
- Fox, D. W., & Price, P. A. 2002, *GRB Circular Network*, 1731, 1
- Fox, D. W., et al. 2003, *ApJ*, 586, L5
- Frail, D. A., et al. 2001, *ApJ*, 562, L55
- Fruchter, A., Krolik, J. H., & Rhoads, J. E. 2001, *ApJ*, 563, 597
- Fruchter, A., Levan, A., Vreeswijk, P., Holland, S. T., & Kouveliotou, C. 2002, *GRB Circular Network*, 1781, 1
- Galama, T. J., et al. 1998, *Nature*, 395, 670
- Galama, T. J., et al. 2000, *ApJ*, 536, 185
- Galama, T. J., et al. 2003, *ApJ*, 587, 135
- Garnavich, P. M., et al. 2003, *ApJ*, 582, 924
- Granot, J., & Sari, R. 2002, *ApJ*, 568, 820
- Granot, J., Nakar, E., & Piran, T. 2003, *Nature*, 426, 138
- Henden, A. 2002, *GRB Circular Network*, 1753, 1
- Holland, S. T., et al. 2004, *AJ*, 128, 1955
- Hjorth, J., et al. 1999, *GRB Circular Network*, 219, 1

- in't Zand, J. J. M., et al. 2001, *ApJ*, 559, 710
- Kinugasa, K., Kato, T., Yamaoka, H., & Torii, K. 2002, *GRB Circular Network*, 1749, 1
- Klose, S., Eisloffel, J., Froebrich, D., Zeh, A., Richter, S., Hoegner, C., & Voss, H. 2002, *GRB Circular Network*, 1739, 1
- Kulkarni, S. R., et al. 1998, *Nature*, 395, 663
- Kumar, P., & Piran, T. 2000, *ApJ*, 535, 152
- Kumar, P., & Panaitescu, A. 2003, *MNRAS*, 346, 905
- Lamb, D. Q., Barentine, J. C., Nysewander, M. C., Reichart, D. E., Schwartz, M., Laws, C., York, D. G., & McMillan, R. J. 2002a, *GRB Circular Network*, 1744, 1
- Lamb, D. Q., Nysewander, M. C., Reichart, D. E., Barentine, J. C., Laws, C., York, D. G., & McMillan, R. J. 2002b, *GRB Circular Network*, 1765, 1
- Lamb, D. Q., et al. 2004, *New Astronomy Review*, 48, 423
- Lazzati, D., Rossi, E., Covino, S., Ghisellini, G., & Malesani, D. 2002, *A&A*, 396, L5
- Lee, B. C., et al. 2001, *ApJ*, 561, 183
- Li, W., Filippenko, A. V., Chornock, R., & Jha, S. 2002, *GRB Circular Network*, 1737, 1
- Li, W., Filippenko, A. V., Chornock, R., & Jha, S. 2003, *ApJ*, 586, L9
- Masetti, N., et al. 2001, *A&A*, 374, 382
- McLeod, B., Caldwell, N., Grav, T., Luhman, K., Garnavich, P., & Stanek, K. Z. 2002, *GRB Circular Network*, 1750, 1
- Moran, J. & Reichart, D. 2004, preprint (astro-ph/0409390)
- Panaitescu, A., & Kumar, P. 2004, *MNRAS*, 353, 511
- Pandey, S. B., Anupama, G. C., Sagar, R., Bhattacharya, D., Castro-Tirado, A. J., Sahu, D. K., Parihar, P., & Prabhhu, T. P. 2003, *A&A*, 408, L21
- Park, H. S., Williams, G., & Barthelmy, S. 2002, *GRB Circular Network*, 1736, 1
- Pravdo, S. H., et al. 1999, *AJ*, 117, 161
- Price, P. A., et al. 2002, *ApJ*, 572, L51

- Price, P. A., et al. 2003, ApJ, 589, 838
- Reichart, D. E. 1999, ApJ, 521, L111
- Reichart, D. E. 2001a, ApJ, 553, 235
- Reichart, D. E. 2001b, ApJ, 554, 643
- Reichart, D. E. 2001c, preprint (astro-ph/0107546)
- Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17
- Sari, R., & Piran, T. 1999, ApJ, 520, 641
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Smith, I. A., et al. 2005, preprint (astro-ph/0503574)
- Stanek, K. Z., et al. 2003, ApJ, 591, L17
- Vreeswijk, P., Burud, I., Fruchter, A., & Levan, A. 2002, GRB Circular Network, 1767, 1
- Waxman, E., & Draine, B. T. 2000, ApJ, 537, 796
- Wei, D. M. 2003, A&A, 402, L9
- Wozniak, P., et al. 2002, GRB Circular Network, 1757, 1
- Zeh, A., Klose, S., & Hartmann, D. H. 2004, ApJ, 609, 952

Table 1. FUN GRB Collaboration Observations of the Afterglow of GRB 021211

Date (UTC)	Mean Δt	Filter	Magnitude ^a	Telescope
Dec 11.4732 ^b	2.84 min	R	15.24 ± 0.07	0.60m Super-LOTIS
Dec 11.4751 ^b	5.63 min	R	16.26 ± 0.12	0.60m Super-LOTIS
Dec 11.4986	39.4 min	I _c	18.60 ± 0.11	0.81m Tenagra II
Dec 11.5114	57.9 min	R _c	19.52 ± 0.13	0.81m Tenagra II
Dec 11.5239	75.9 min	V	20.06 ± 0.41	0.81m Tenagra II
Dec 11.5366	94.1 min	B	>19.8	0.81m Tenagra II
Dec 11.5479	1.84 hr	I _c	19.99 ± 0.24	0.81m Tenagra II
Dec 11.5525	1.95 hr	R _c	20.74 ± 0.42	0.81m Tenagra II
Dec 11.5566	2.05 hr	R _c	20.70 ± 0.16	0.65m Gunma
Dec 11.9583	11.7 hr	R _c	>22.0	1.34m Tautenburg
Dec 11.9744	12.1 hr	I _c	>20.7	1.34m Tautenburg
Dec 12.3883	22.0 hr	i*	23.02 ± 0.12	3.5m ARC
Dec 13.4680	47.9 hr	V	23.0 ± 0.5	1.0m USNO
Dec 28.4283	17.0 day	i*	24.41 ± 0.22	3.5m ARC
Mar 23.1335	102 day	i*	24.51 ± 0.29	3.5m ARC

^aUpper limits are 3σ .

^bFlux weighted using an iterated power-law index of $\alpha = -1.37$ (§2.1).

Fig. 1.— i^* light curve from 22 hours to 102 days after the burst and best-fit WIND-BLUE model from §3.1. The host galaxy dominates at late times. We do not detect the supernova (Fruchter et al. 2002; Della Valle et al. 2003), likely due to the timing of our observations.

Fig. 2.— *Top panel:* BVRR_cI_cJHK_s light curves from 2.8 min to 2.0 days after the burst and best-fit WIND-BLUE model from §3.1. Upper limits are 3σ . We do not include the dotted K_s upper limit in our fits. Data are from McLeod et al. (2002), Pandey et al. (2003), Fox et al. (2003), Holland et al. (2004), and this paper. *Bottom panel:* $g'r'i^*$ and unfiltered KAIT and NEAT light curves from 9.2 min to 1.0 days after the burst and best-fit WIND-BLUE model from §3.1. The dotted curves are the reverse and forward shock components of the best-fit model for the spectral response of KAIT’s CCD. Data are from Fox et al. (2003), Li et al. (2003), and this paper.

Fig. 3.— Residuals of Figure 2. Colors are the same as in Figure 2. Notice the increase relative to the best-fit model of the unfiltered NEAT and KAIT data, which is also clearly visible in Figure 2b, concurrent with a decrease relative to the best-fit model of our R_c and possibly I_c data from Tenagra and Gunma. The dashed curves are our best-fit simple model for this from §3.2.

Fig. 4.— Spectral flux distribution at six epochs and best-fit WIND-BLUE model from §3.1 (solid curves). Dashed curves are the same fit, but with source-frame extinction set to zero. We scale data to these times using the best-fit light curve and when there are multiple points per spectral band we plot weighted averages of the scaled data for clarity (see §3.2). Colors are the same as in Figure 2. Horizontal bars mark the 90% width of the filter. Upper limits are 3σ . We do not include the dotted K_s upper limit in our fits.









